



Folio B is an organic foliar fertilizer used in the absence of boron. Folio B is used after a leaf analysis or as a preventative before periods of increasing mineral demand. Such periods are during flowering, fruiting or undergoing strong growth and development.

Folio B contains a high concentration of boron in combination with a dispersant. The dispersant ensures that the mineral is in good contact with the leaf surface and the growth has a good opportunity to absorb it. The dispersant is made from plant material and is organically transferable.

Finished formulated foliar fertilizer for use in drilling deficiency by spraying foliage in vegetables, fruits, legumes and grains. Enhances plant development of buds and leaves. Stimulates the flower, formation and growth of the fruit.

To best help the growths out of a mineral deficiency, Folio B is added energy in the form of a foliar fertilizer (Pepton). Folio B is absorbed in the growth within a few hours and the effect may be visible already on the same day.

Folio B is supplied in powder form and should be mixed with water and can be used with other types of foliar fertilizers or pesticides. Folio B is dosed at between 50 and 200 grams (dry weight) per day. acres. Mix in water according to desired dosage, experience, soil or leaf analysis. Fertilizer approved in organic farming by (EC) No 834/2007.

<p>Apple Strengthens growth and development: Before flowering at risk of deficiency, otherwise after flowering is complete. Can be repeated after 10-14 days, but not after leaves are fallen.</p>	<p>Potato Strengthens growth and development: Used one week after full germination, repeat once after 14 days, etc. Repeat up to 2-3 weeks after tuber formation.</p>	<p>Lettuce in open air Strengthens growth and development: 10-14 days after planting/germination. In case of moderate deficiency repeat at 10-14 day intervals.</p>	<p>Cauliflower/Cabbage Stimulate initial head formation. Especially at low soil temperatures in spring / early summer. From the 4-leaf stage. Repeat at 10 to 14 day intervals for moderate deficiency symptoms.</p>	<p>Raspberry First treatment when the buds are green, second when they are white. The plants can also be treated after harvest.</p>
---	--	--	---	--



Analysis

Boron content (B)	min. 11,2 %
Pepton	20 %
Organic carbon	ca. 4 %



Content

All amino acids (in the Pepton part) are of type L	
alanine	threonine
lysine	glycine
arginine	tryphophan
methionine	histidine
aspartic acid	tyrosine
phenylalanine	isoleucine
cysteine	valine
serine	leucine
glutamic acid	proline



Recommendation

Dissolves in water:
50 - 200 grams (dry weight) per decar

10 kg