



Folio Cu is an organic foliar fertilizer that is used after a leaf analysis or as a preventative before periods of increasing mineral demand. Copper (Cu) is one of the eight essential micronutrients. It is needed by plants in small quantities, but still important in a wide range of development processes.

The availability of copper depends on soil type. Folio Cu contains a high concentration of copper in combination with a dispersant. Finished formulated foliar fertilizer for use against copper deficiency by spraying on foliage in vegetables, fruits, legumes, grains and oilseeds. The dispersant ensures that the mineral is in good contact with the leaf surface and the growth has a good opportunity to absorb it. The dispersant is made from plant material and is organically transferable. To best help the growth out of a mineral deficiency, Folio Cu is added energy in the form of a foliar fertilizer (Pepton).

Folio Cu is absorbed in the growth within a few hours and the effect may be visible already on the same day. Strengthens the plant's development of stems, buds and leaves. Stimulates the flower, formation and growth of the fruit.

Folio Cu is supplied in powder form and should be mixed with water and can be used with other types of foliar fertilizers or pesticides.

Mix in water according to desired dosage, experience, soil or leaf analysis. Fertilizer approved in organic farming by (EC) No 834/2007.

- | | | | | |
|--|--|--|---|--|
| <p>Carrot
Strengthens growth and development: When the plants are 15 cm high. In case of moderate deficiency, treatment is repeated at 10-14 day intervals.</p> | <p>Cauliflower
Strengthens growth and development: At 4 to 6 leaf stage. For Moderate copper deficiency should be repeated once or twice every 10 to 14 days.</p> | <p>Lettuce in open air
Strengthens growth and development: 10-14 days after planting / germination. In case of moderate deficiency repeat with 10-14 days interval. Last treatment at least 1 month before harvest.</p> | <p>Grain
Strengthens growth and development: From the two-leaf stage to the first joint node becomes visible. By moderate deficiency is repeated at 10 to 14 day intervals. In autumn wheat treatment before winter.</p> | <p>Raspberry / Strawberry
Strengthens growth and development: Early in the season, ie before flowering. May also be use after season.</p> |
|--|--|--|---|--|



Analysis

Copper content (Cu)	min. 12 %
Pepton	20 %
Organic carbon	ca. 4 %



Content

All amino acids (in the Pepton part) are of type L

alanine	threonine
lysine	glycine
arginine	tryphophan
methionine	histidine
aspartic acid	tyrosine
phenylalanine	isoleucine
cysteine	valine
serine	leucine
glutamic acid	proline



Recommendation

Dissolves in water:
50 - 200 grams (dry weight) per acre.

10 kg