



grønn
gjødsel



Folio Ca is a pre-formulated foliar fertilizer for use against calcium deficiency in potato, fruit and berries. Used after leaf analysis or as a preventative before periods of increasing mineral demand. Such periods are flowering, fruiting or strong growth and development.

Folio Ca contains a high concentration of calcium in combination with a dispersant. The dispersant ensures that the mineral is in good contact with the leaf surface and the growth has a good opportunity to absorb it. The dispersant is made from plant material and is organically transferable. To best help the growths out of a mineral deficiency, Folio Ca is added energy in the form of a foliar fertilizer (Pepton). Folio Ca is absorbed into the growths within a few hours and the effect may be visible already

on the same day. Folio Ca is supplied in powder form and should be mixed with water and can be used with other types of foliar fertilizers or pesticides. Folio is dosed at between 50 and 200 grams (dry weight) per day. acres. Enhances plant development of buds and leaves. Stimulates the flower, formation and growth of the fruit. Mix in water according to desired dosage, experience, soil or leaf analysis. Folio Ca is an ecologically approved corrective for use in organic farming by (EC) No 834/2007.

Apple

For development and color: 3-8 treatments at 10-14 day intervals from petal drop. For better color of the apples treatment when the fruit has begun to change color, often repeated after 7 days, normally 2-3 weeks for harvest.

Potato

Stimulate early growth (1), increase tuber (2) and size (3): (1) Used one week after germination. (2) Used when half of the chair tops are twice as thick as the rest. (3) Two treatments when the tubers are about 10 mm, repeat once after 14 days.

Lettuce in open air

Strengthens growth and development: 2-3 treatments starting 10-14 days after planting or germination. Treatment is repeated at 7-10 day intervals.

Cauliflower

Stimulate initial head formation, especially at low soil spring / early summer. 2-3 treatments at the point of growth to first head formation. Especially at low soil temperatures in spring / early summer. Repeats every 7 to 14 days.

Stone fruit

Stimulate growth and development: 2-5 treatments at 7-14 day intervals from petal drop.



Analysis

Calcium content (Ca)	min. 18 %
Peptone	20 %
Organic carbon	ca. 4 %



Content

All amino acids (in the Pepton part) are of type L

alanine	threonine
lysine	glycine
arginine	tryphophan
methionine	histidine
aspartic acid	tyrosine
phenylalanine	isoleucine
cysteine	valine
serine	leucine
glutamic acid	proline



Recommendation

Dissolves in water:
50 - 200 grams (dry weight) per decare

10 kg