



Ready-formulated foliar fertilizer for use against iron deficiency for spraying on foliage in vegetables, fruits, legumes and grains. Strengthens the plant's development of buds and leaves. Stimulates the flower, the formation and growth of the fruit.

Folio Fe is used after leaf analysis or prophylactically before periods of increasing mineral demand. Such periods are during flowering, fruiting or during strong growth and development. Folio Fe contains a high concentration of iron in combination with a dispersant. The dispersant ensures that the mineral has good contact with the leaf surface, and the growth has a good opportunity to absorb it. The dispersant is made from plant material and is organically marketable. To best help the growth out of a mineral

deficiency, Folio Fe is added energy in the form of a foliar fertilizer (Pepton). Folio Fe is absorbed into the growths within a few hours and the effect can be visible already on the same day. Folio Fe is supplied in powder form and should be mixed with water, and can be used with other types of foliar fertilizers or pesticides. Folio Fe is dosed with between 50 and 200 grams (dry weight) per. acres. Folio Fe is an organically approved corrective for use in organic farming by (EC) no. 834/2007.

Apple

Strengthens growth and development: Before flowering at risk of deficiency, otherwise after flowering is complete. Can be repeated after 10-14 days. For cork crust exposed types, treatment is usually postponed until 6 weeks laterflowering.

Potato

Strengthens growth and development: Use one week after full germination, repeat once after 14 days, etc.

Salad open air

Strengthens growth and development: 10-14 days after planting/germination. In case of moderate deficiency repeat at 10-14 day intervals. Last treatment no later than 1 month before harvest.

Cauliflower

Stimulates incipient head formation, especially at low soil temperature spring / early summer. From the 4-leaf stage. Repeat at 10 to 14 day intervals for moderate deficiency symptoms.

Legumes

Strengthens growth and development: Start at 4-6 leaf stage to prevent general deficiency. At the start of flowering, repeat at the end of flowering to prevent internal discoloration.



Analysis

Iron content (Fe)
Pepton
Organic carbon

ca. 19 %
20 %
ca. 4 %



Content

All amino acids are of type L:

Alanin
Lysin
Arginin
Metionin
Asparaginsyre
Fenylalanin
Cystein
Serin
Glutaminsyre

Treonin
Glycin
Tryptofan
Histidin
Tyrosin
Isoleucin
Valin
Leucin
Prolin

10 kg



Recommendation

Mix in water according to desired dosage, according to experience, soil or leaf analysis

50 - 200 grams (dry weight) per. acre.